

Causality Violation in Non-local Quantum Field Theory

Ambar Jain and Satish D. Joglekar

1st February 2008

Department of Physics, I.I.T.Kanpur, Kanpur 208016 (INDIA)

Abstract

We study the causality violation in the non-local ϕ^4 -theory (as formulated by Kleppe and Woodard) containing a finite mass scale Λ . Starting from the Bogoliubov-Shirkov criterion for causality, we construct and study combinations of S-matrix elements that signal violation of causality in the one loop approximation. We find that the causality violation in the exclusive process $\phi + \phi \rightarrow \phi + \phi$ grows with energy, but the growth with energy, (for low to moderate energies) is suppressed to all orders compared to what one would expect purely from dimensional considerations. We however find that the causality violation in other processes such as $\phi + \phi \rightarrow \phi + \phi + \phi + \phi$ grows with energy as expected from dimensional considerations at low to moderate energies. For high enough energies comparable to the mass scale Λ , however, we find a rapid (exponential-like) growth in the degree of causality violation. We suggest a scenario, based on an earlier work, that will enable one to evade a large theoretical causality violation at high energies, should it be unobserved experimentally.

1 INTRODUCTION

The chief problem one is faced with any local quantum field theory (LQFT) is the presence of ultra-violet divergences. A number of regularization procedures have been proposed to deal with such divergences. The basic problem of divergences arises from the ill-defined nature of a product of two local field operators at the same space-time point [1] in a LQFT. One of the natural suggestions to deal with this problem therefore has been that we should attempt to replace a local interaction by a nonlocal (smeared) interaction which somehow would take care of the divergences arising from two local field operators at the same space-time point [2]. While making such an attempt, however, there is a danger that one may lose physically meaningful properties of the local theory such as unitarity, stability [3, 4], gauge invariance (in the case of gauge theories) and causality [5]. Many attempts have been made in this direction to construct a finite and physically meaningful quantum field theory with the use of nonlocal interactions [2, 5]. One of the physical requirements of the physical theory is the unitarity of the S-Matrix. This requires that in the process of introducing nonlocality, one does not introduce any unphysical ghost degrees of freedom that can contribute to an S-matrix. Also, for local theories with a local gauge-invariance, it is necessary to introduce nonlocal interactions in a manner that is compatible with the local gauge symmetry (or an equivalent non-local symmetry), so that the essential consequences of a local gauge symmetry are preserved after non-localization.

An elegant formulation of nonlocal quantum field theory that preserves unitarity and (an equivalent of) local gauge symmetry has been given by Kleppe and Woodard [5]. It was based upon the earlier works by Moffat[6], Elizer and Woodard [3, 4] and on an explicit construction for QED by Evens et al [7]. A systematic procedure for constructing a non-local action from a local QFT

has been presented in Ref. [5]. Equally importantly, this work has established that the local/global symmetries can be preserved in their nonlocal form and the WT identities of a local QFT derivable from local symmetries such as gauge invariance/BRS symmetry find their natural nonlocal extensions (at least in the Feynman gauge). It has since been studied extensively [8, 9]. It has also been found useful for regularizing theories with supersymmetry [10] (where dimensional regularization does not work) and for the regularization of BV theories [11].

This formulation of a non-local field theory can be looked upon either as a regularization [7, 8] or as a physical theory with a finite mass parameter Λ [6, 5]. In either case, and for *any finite* Λ , the theory is unitary and has an appropriate generalization of a local gauge symmetry. The latter of these interpretations can further be looked upon in a number of distinct ways. One could think the non-locality as representing a form factor with a momentum cut-off Λ [6]. One could also think of this theory as embodying a granularity of space-time of the scale $1/\Lambda$ or as an intrinsic mass scale Λ [5, 12, 14, 13]. One could also consider such a theory as representing an effective field theory valid when the energy scale involved is smaller than Λ [15].

In an earlier work [14], we have found such a non-local formulation *with a finite* Λ , very useful in understanding the renormalization program in the renormalizable field theories. We have shown that this formulation enables one to construct a mathematically consistent framework in which the renormalization program can be understood in a natural manner. The framework does not require any violations of mathematical rigor usually associated with the renormalization program. This framework, moreover, made it possible to theoretically estimate the mass scale Λ . The nonlocal formulations can also be understood [15] as an effective field theory formulation of a physical theory that

is valid up to mass scale $\sim \Lambda$. In such a case, the unknown physics at energy scales higher than Λ [such as a structure in terms of finer constituents, additional particles, forces, supersymmetry etc] can *effectively be represented* in a *consistent* way (a unitary gauge-invariant finite (or renormalizable) theory) by the non-local theory. In other words, the nonlocal standard model can serve as such an effective field theory [15] and will afford a model-independent way of reparametrizing the effects beyond standard model consistently.

Despite the various advantages of these nonlocal formulations, it suffers from some limitations: such a theory, while it preserves causality at the tree-level S-matrix (which is the *same as* the local one), has quantum violations of causality, expected to be small by a factor $\sim \frac{g^2}{16\pi^2}$. In this work, we shall carry out an elementary study of the question of causality violation (CV) in NLQFT . We shall adopt the approach of Bogoliubov and Shirkov [17], and use Bogoliubov-Shirkov criterion [18] to study CV. Bogoliubov-Shirkov criterion (See section 2.3) allows one obtain certain combinations of S-matrix elements that must vanish if causality is preserved; and therefore these combinations constitute quantities that can characterize the CV in a NLQFT. We study these in the 1-loop approximation for the $\lambda\phi^4$ theory, this being the simplest field theory model where definite conclusions can be drawn.

We shall now introduce the plan of the paper. In section 2, we shall introduce the formalism of non-local field theories for the $\lambda\phi^4$ theory and give Feynman rules. We shall also summarize the causality condition as formulated by Bogoliubov and Shirkov. We shall construct quantities that can be used to characterize the causality violation. We shall also make brief comments on the possible violation of causality in LQFT [19] and its distinction from the causality violation we shall address to in the present context of the NLQFT. We shall also summarize the scenario suggested in [15]. In Section 3, we shall eval-

uate the causality violation effects at the one loop level in two of the exclusive processes: (i) 2 scalars $\rightarrow$ 2 scalars, (ii) 2 scalars $\rightarrow$ 4 scalars. We shall show that at an energy scale small compared to Λ , the causality violation in the first process is of an order smaller than what would be expected from purely dimensional consideration. We shall further show that at energies comparable to Λ or greater, we shall find an exponential-like growth in the causality violation. For the second process, we find that we have a causality violation as expected from general dimensional arguments for energy scales $\ll \Lambda$. On the other hand, at energy scales $\sim \Lambda$ we again find an exponential-like growth. In section 4, we shall give a general argument to show that the results noted in the one loop for the 2 scalars $\rightarrow$ 2 scalars ought to be valid to all orders. In section 5, we shall present a scenario based on that presented in [15], that allows one to evade a large scale causality violation should it be unobserved experimentally.

2 PRELIMINARIES

In this section, we shall summarize, for our future use, various known results on the non-local field theories, Bogoliubov-Shirkov criterion of causality, and brief comments on possible causality violation in LQFT and the pertinence (or its lack) of these results to the present discussion. We will also summarize the view-point regarding the non-local field theories as presented in [15].

2.1 NON-LOCAL REGULARIZATION:

We shall first review, very briefly, the construction of a non-local field theory with a finite mass scale Λ , given its local counterpart. We shall present the construction with reference to the $\lambda\phi^4$ theory as presented in [5]; since that suffices for our purpose.

We start with the local action for a field theory, in terms of a generic field

ϕ , as the sum of the quadratic and the interaction part:

$$S[\phi] = F[\phi] + I[\phi]$$

and express the quadratic piece as

$$F[\phi] = \int d^4x \phi_i(x) \mathfrak{S}_{ij} \phi_j(x)$$

We define the regularized action in terms of the smeared field $\hat{\phi}$, defined in terms of the kinetic energy operator $\mathfrak{S}_{ij}$ as,

$$\hat{\phi} \equiv \mathcal{E}^{-1} \phi \quad \mathcal{E} \equiv \exp[\mathfrak{S}/\Lambda^2]$$

The nonlocally regularized action is constructed by first introducing an auxiliary action $S[\phi, \psi]$. It is given by

$$S[\phi, \psi] = F[\hat{\phi}] - A[\psi] + I[\phi + \psi]$$

where ψ is called a “shadow field” with an action

$$A[\psi] = \int d^4x \psi_i O_{ij}^{-1} \psi_j$$

with O defined by

$$O \equiv \frac{\mathcal{E}^2 - 1}{\mathfrak{S}}$$

The action of the non-local theory is defined as

$$\hat{S}[\phi] = S[\phi, \psi] \big|_{\psi=\psi[\phi]}$$

where $\psi[\phi]$ is the solution of the classical equation

$$\frac{\delta S}{\delta \psi} = 0$$

The Feynman rules for the nonlocal theory are simple extensions of the local ones. The vertices are unchanged but every leg can connect either to a smeared propagator

$$\frac{i\mathcal{E}^2}{\Im + i\epsilon} = -i \int_1^\infty \frac{d\tau}{\Lambda^2} \exp\left\{\frac{\Im\tau}{\Lambda^2}\right\}$$

or to a shadow propagator [shown by a line crossed by a bar]

$$\frac{i[1 - \mathcal{E}^2]}{\Im + i\epsilon} = -iO = -i \int_0^1 \frac{d\tau}{\Lambda^2} \exp\left\{\frac{\Im\tau}{\Lambda^2}\right\}$$

In the context of the $\lambda\phi^4$ theory, we have,

$$\Im = -\partial^2 - m^2 \quad I(\phi) = -\frac{\lambda}{4}\phi^4$$

The Feynman rules for propagator in momentum space read:

1. For the ϕ -propagator (smeared propagator) denoted by a straight line:

$$i \frac{\left\{ \exp\left[\frac{p^2 - m^2}{\Lambda^2}\right] \right\}}{p^2 - m^2 + i\epsilon}$$
2. For the ψ -propagator denoted by a *barred* line:

$$i \frac{\left\{ 1 - \exp\left[\frac{p^2 - m^2}{\Lambda^2}\right] \right\}}{p^2 - m^2 + i\epsilon}$$

3. The 4-point vertex is as in the local theory, except that any of the lines emerging from it can be of either type. (There is accordingly a statistical factor).
4. In a Feynman diagram, the internal lines can be either shadow or smeared,

with the exception that no diagrams can have closed shadow loops.

2.2 Condition of Causality

The causality condition that we have used to investigate causality violation in NLQFT is the one discussed by Bogoliubov and Shirkov [17]. They have shown that an S-matrix for a theory that preserves causality must satisfy the condition

$$\frac{\delta}{\delta g(x)} \left(\frac{\delta S[g]}{\delta g(y)} S^\dagger[g] \right) = 0 \quad \text{for } x < \sim y \quad (1)$$

where $x < \sim y$ means that either $x^0 < y^0$ or x and y are space like separated. The above condition has been formulated treating the coupling $g(x)$ as space-time dependent. This is used together with the unitarity condition

$$S^\dagger[g]S[g] = 1 \quad (2)$$

to obtain the form of the S-matrix in LQFT. We shall use them together to study the causality violation in NLQFT. To extract the physics from the above mentioned causality condition, in the perturbative sense, S-matrix is written as a functional Taylor expansion in powers of coupling $g(x)$:

$$S[g] = 1 + \sum_{n \geq 1} \frac{1}{n!} \int S_n(x_1, \dots, x_n) g(x_1) \dots g(x_n) dx_1 \dots dx_n, \quad (3)$$

in which $S_n(x_1, \dots, x_n)$ are *operator* expressions (symmetric in all arguments) which depend upon the field operators and on their partial derivatives at the points $x_1, \dots, x_n$. We note that experimental information is not however about an $S_n(x_1, \dots, x_n)$ but about

$$S_n \equiv \int S_n(x_1, \dots, x_n) dx_1 \dots dx_n$$

obtained by putting $g(x) = \text{constant} = g$. As shown in [18], one can obtain perturbatively useful causality conditions for each $n = 1, 2, 3, \dots$ from (1) and (2) and using the expansion (3):

$$H_n(y, x_1, \dots, x_n) = iS_{n+1}(y, x_1, \dots, x_n) + i \sum_{0 \leq k \leq n-1} P\left(\frac{x_1, \dots, x_k}{x_{k+1}, \dots, x_n}\right) S_{k+1}(y, x_1, \dots, x_k) S_{n-k}^\dagger(x_{k+1}, \dots, x_n) = 0 \quad (4)$$

Here, $P\left(\frac{x_1, \dots, x_k}{x_{k+1}, \dots, x_n}\right)$ is defined as the sum over the distinct ways of partitioning $(\frac{n!}{k!(n-k)!})$ in number $\{x_1, x_2, x_3, \dots, x_n\}$ into two sets of k and $(n-k)$ (such as $\{x_1, x_2, x_3, \dots, x_k\} \{x_{k+1}, \dots, x_n\}$). Similarly perturbatively useful unitarity condition, for each n , is given by

$$S_n(x_1, \dots, x_n) + S_n^\dagger(x_1, \dots, x_n) + \sum_{1 \leq k \leq n-1} P\left(\frac{x_1, \dots, x_k}{x_{k+1}, \dots, x_n}\right) S_k(x_1, \dots, x_k) S_{n-k}^\dagger(x_{k+1}, \dots, x_n) = 0.$$

For $n = 1, 2$ respectively, causality condition is explicitly given by

$$H_1(x, y) \equiv iS_2(x, y) + iS_1(x)S_1^\dagger(y) = 0 \quad (5)$$

$$H_2(x, y, z) \equiv iS_3(x, y, z) + iS_1(x)S_2^\dagger(y, z) + iS_2(x, y)S_1^\dagger(z) + iS_2(x, z)S_1^\dagger(y) = 0 \quad (6)$$

and unitarity condition is given by

$$S_1(x) + S_1^\dagger(x) = 0 \quad (7)$$

$$S_2(x, y) + S_2^\dagger(x, y) + S_1(x)S_1^\dagger(y) + S_1(y)S_1^\dagger(x) = 0 \quad (8)$$

The unitarity conditions could be used with causality condition to replace $S_n^\dagger$ factor in (4) by appropriate S_m terms. One can then evaluate the matrix el-

elements of $H_n(y, x_1, \dots, x_n)$ between appropriate initial and the final states to convert the relations (4) in terms of the S-matrix *element coefficients* in (3). In the case of the local theory, these relations are trivially satisfied. In the case of the nonlocal theories, these quantities, on the other hand, afford a way of characterizing the causality violation. Further, these quantities contain however not the usual S-matrix elements that one can observe in an experiment (which are obtained with a *constant i.e. space-time-independent* coupling), but rather the coefficients in (3). We thus find it profitable to construct appropriate space-time integrated versions out of $H_n(y, x_1, \dots, x_n)$. Thus, for example, we can consider

$$H_1 \equiv \int d^4x \int d^4y [\vartheta(x_0 - y_0)H_1(x, y) + \vartheta(y_0 - x_0)H_1(y, x)] \quad (9)$$

The causality condition (5) would then imply that a lack of CV requires that¹ $H_1 \equiv 0$. Conversely, a non-zero H_1 necessarily implies CV. Recalling that $S_2(x, y) = S_2(y, x)$, we can write

$$H_1 = i \int d^4x \int d^4y S_2(x, y) - i \int d^4x \int d^4y T[S_1(x)S_1(y)]$$

which can be expressed entirely in terms of Feynman diagrams that appear in the usual S-matrix amplitudes. In a similar manner, we can formulate

$$H_2 \equiv \int d^4x \int d^4y \int d^4z H_2(x, y, z) \vartheta(x_0 - y_0) \vartheta(y_0 - z_0) \quad (10)$$

$$+ 5 \text{ symmetric terms} \quad (11)$$

and can itself be expressed in terms of Feynman diagrams.

Finally, we comment on the counterterms in the above equations. When we wrote the original causality condition in terms of $H_1(x, y)$, with $y < \sim x$,

¹These statements are subject to the counterterms in the definition of H_1 to be discussed shortly.

x and y were distinct points and there were no divergences in both the terms that are present in $H_1(x, y)$. On the other hand, when we convert it to H_1 defined above, (each term in its definition now allows for coincident points), and contains a "subtraction" or a counterterm. The counterterm² *present* in $\int d^4x \int d^4y S_2(x, y)$ makes this term finite while the "subtraction" that *has* to be included in $\int d^4x \int d^4y T[S_1(x)S_1(y)]$ has to be such that as $\Lambda \rightarrow \infty$ (i.e. the local limit) H_1 must vanish. This criterion, however, does not uniquely determine the net counterterm in H_1 ; there is an ambiguity of a *constant, i.e. momentum-independent* term that vanishes as $\Lambda \rightarrow \infty$. We shall assume, for definiteness, that experiments show no causality violation at low energies, and we shall adjust this counterterm accordingly. Nonetheless, the presence of a *momentum-dependent* CV terms in H_1 necessarily signals CV at *some* energy scale, irrespective of the ambiguity in the counterterm.

2.3 Causality Violation in Local Quantum Field Theories

We shall briefly discuss the issue of causality violation in the LQFT and point out the distinction between the issue there and here in the context of the NLQFT. There is much discussion in literature about the CV in LQFT and is thought to be still a controversial matter [19]. However, these discussions revolve around processes that take place in a *finite duration*, unlike an S-matrix which relates to the time-evolution of a scattering system over an infinite duration. The possible signal of CV in LQFT in fact vanishes as the time of observation of a process goes to infinity. This is in fact consistent with no CV in LQFT according to the BS criterion; which refers to an S-matrix element.

In light of Bogoliubov and Shirkov formulation of S-matrix theory [18] and construction of S_n , which is widely used in perturbative QFT, it is clear that

²The expansion of (3) is the expansion of the *renormalized* S-matrix in terms of the *renormalized* coupling g .

causality violation does not occur in the S-matrix theory in a LQFT. Therefore we have studied CV in NLQFT in the S-matrix formalism.

2.4 Non-local Field Theory as an Effective Field Theory

In this subsection, we shall summarize the view-point presented in references [15] and [14] regarding the non-local theories as effective field theories valid up to a mass scale Λ . The basic idea presented in [15] was with reference to a quantum field theory for which the LSZ formulation leads to the condition $0 < Z < 1$. In a local quantum field theory, this condition is ignored in perturbation theory as the wave-function renormalization Z diverges. It was suggested in [15] that we can resurrect this condition in a NLQFT with an intrinsic scale Λ , and in fact give a meaning to the scale and make an estimate using the condition $0 < Z < 1$ *as applied in perturbation theory*. The idea was to require that the mass-scale Λ (together with the coupling) be such that the calculated Z satisfies $0 < Z < 1$. This lead to a condition of the type

$$\frac{g^2}{16\pi^2} \ln \frac{\Lambda^2}{m^2} \lesssim 1.$$

The suggested interpretation of the NLQFT then was (i) It is an effective theory valid up to energy scale Λ ;(ii) For energy scales beyond Λ , one needs to replace the NLQFT by a more fundamental theory (of constituents) having its own (larger) mass scale Λ' and a coupling g' and such constituent fields that the condition $0 < Z' < 1$ is satisfied again within the energy domain $\Lambda < E \lesssim \Lambda'$.

An equivalent discussion can be given even for theories for which such a condition is not available. It is based on the work [14], in which we have attempted to give a mathematically rigorous understanding of the renormalization program using the framework of the NLQFT. There, we were lead to a very similar condition if the renormalization program is to be understood in a rigorous way.

We can then impose exactly similar interpretation on the NLQFT.

3 Causality Violation at One Loop

3.1 for 2 Particle $\rightarrow$ 2 Particle

We shall now consider the contributions to H_1 , of (9) a quantity quantifying the causality violation in the theory, for the above process. There are two contributions to this term: (a) One set of contributions comes from the diagrams with a shadow propagator (see fig. 1(a)) and analogous diagrams in t - and the u - channels. These contribute to $\int d^4x \int d^4y S_2(x, y)$ but not to the other term. (b) The second contribution comes from the renormalization counterterms present in $\int d^4x \int d^4y S_2(x, y)$ and $\int d^4x \int d^4y T[S_1(x)S_1(y)]$. The counterterm is (partly) determined by the requirement that the causality violation vanishes as $\Lambda \rightarrow \infty$.

(a) We have evaluated these diagrams in the massless limit and for non-zero mass.




Figure 1: (a) 2 particle $\rightarrow$ 2 particle, s-channel diagram contributing to CV
 (b) 2 particle $\rightarrow$ 4 particle, a sample diagram contributing to CV

3.1.1 Massless Limit

After some involved calculations, we find that the amplitude for s-channel diagram in the massless limit is given by

$$\Gamma_0(s) = \frac{9\lambda^2}{4\pi^2} \sum_{n=0}^{\infty} \frac{\left(\frac{s}{\Lambda^2}\right)^n \left(1 - \frac{1}{2^n}\right)}{n((n+1)!)}.$$
 (12)

The above expression, a convergent series, is particularly useful when $s \ll \Lambda^2$. In addition, there are the t -channel and the u -channel diagrams each of which is given by an expression (12) with $s \rightarrow t$ and $s \rightarrow u$ respectively. The sum of three such diagrams, for the case $s \ll \Lambda^2$ is given by,

$$\begin{aligned} \Gamma_0(s) + \Gamma_0(t) + \Gamma_0(u) &= \frac{9\lambda^2}{4\pi^2} \left\{ 3\ln 2 + \frac{1}{4\Lambda^2}(s+t+u) + O\left(\frac{s^2}{\Lambda^4}\right) \right\} \quad (13) \\ &= \frac{9\lambda^2}{4\pi^2} \left\{ 3\ln 2 + \frac{1}{\Lambda^2}(m^2) + O\left(\frac{s^2}{\Lambda^4}, \frac{t^2}{\Lambda^4}, \frac{u^2}{\Lambda^4}\right) \right\} \quad (14) \end{aligned}$$

The total causality violating amplitude is thus given by sum of the above result and the renormalization counterterm. One effect of the latter is to subtract out from

$$\Gamma_0(s) + \Gamma_0(t) + \Gamma_0(u)$$

the contribution that does not vanish away as $\Lambda \rightarrow \infty$. As explained in the section 2.2, we shall fix the counterterm further by assuming that CV vanishes at low momenta³ ($s = t = u = 0$). We thus obtain the net causality violating amplitude:

$$\frac{9\lambda^2}{64\pi^2} \left(\frac{s^2 + t^2 + u^2}{\Lambda^4} \right).$$
 (15)

and thus is a *quadratic* function of the basic momentum variables s, t and u . Consequently, it grows much more slowly than expected with energy at low energies. One would therefore have to go to much higher energies to see causality

³Please recall the comment in section 2.2.

violation than one would naively expect on the basis of dimensional arguments. In the section 4, we shall give a general argument to show that this must be so to all orders.

We shall now study the result for energies comparable to Λ . To see the behavior, we focus our attention on $\Gamma_0(s)$. We note that the series,

$$\Gamma_0(s) = \frac{9\lambda^2}{4\pi^2} \sum_{n=0}^{\infty} \frac{\left(\frac{s}{\Lambda^2}\right)^n \left(1 - \frac{1}{2^n}\right)}{n((n+1)!)}.$$
 (16)

for $s \sim \Lambda^2$ grows exponentially. To see this, we introduce

$$A(\xi) \equiv \sum_{n=1}^{\infty} \frac{\xi^n}{n(n+1)!}$$

We can then easily verify the following bounds for any $\xi > 0$,

$$\left(\frac{e^\xi}{\xi^2} - \frac{1}{\xi^2} - \frac{1}{\xi} - 1/2\right) < A(\xi) < \frac{1}{9} \left(\frac{e^{3\xi}}{\xi^2} - \frac{1}{\xi^2} - \frac{3}{\xi} - 9/2\right)$$

[The lower bound is sufficient to conclude an exponential growth. The upper bound is a rather loose bound and can be made stricter, but that is not required].

We thus find that as $s \gtrsim \Lambda^2$, there is an exponentially rising causality violation from the s -channel diagram.

3.1.2 Non-zero Mass

For completeness, we shall briefly discuss the case of $m \neq 0$; even though it does not lead to any substantial changes in conclusions. After invoking Schwinger parameterization and some calculations, we obtain the s -channel amplitude for this case in the following integral form:

$$\Gamma(s) \equiv \frac{9\lambda^2}{4\pi^2} \int_0^{1/2} dx \int_{\frac{1}{1-x}}^{\frac{1}{x}} \frac{d\tau}{\tau} \exp\left\{-\frac{\tau}{\Lambda^2} (m^2 - x(1-x)s)\right\}. \quad (17)$$

Expanding in powers of s we get from the above expression

$$\Gamma(s) = \sum_{n=0}^{\infty} \gamma_n(m, \Lambda) s^n, \quad (18)$$

where first three γ_n are given by

$$\gamma_0(m, \Lambda) = \frac{9\lambda^2}{4\pi^2} \int_0^{1/2} dx \int_{\frac{1}{1-x}}^{1/x} \frac{d\tau}{\tau} e^{-\frac{m^2\tau}{\Lambda^2}}, \quad (19)$$

$$\gamma_1(m, \Lambda) = \frac{9\lambda^2}{4\pi^2} \int_0^{1/2} dx \int_{\frac{1}{1-x}}^{1/x} d\tau \frac{x(1-x)}{\Lambda^2} e^{-\frac{m^2\tau}{\Lambda^2}}, \quad (20)$$

$$\gamma_2(m, \Lambda) = \frac{9\lambda^2}{4\pi^2} \int_0^{1/2} dx \int_{\frac{1}{1-x}}^{1/x} \tau d\tau \frac{x^2(1-x)^2}{2\Lambda^4} e^{-\frac{m^2\tau}{\Lambda^2}}. \quad (21)$$

The contributions to the total amplitude from the $n = 0$ term is cancelled by the renormalization counterterm. Since $s + t + u = 4m^2$, the contribution from the $n = 1$ terms add up to a momentum independent term. $\gamma_2(m, \Lambda)$ could be further simplified in terms of incomplete Γ -functions but the expression is lengthy and complicated. To the lowest order it has term $(const.)s^2/\Lambda^4$ and at next leading order there are terms like m^2s^2/Λ^6 and $\ln(m^2/\Lambda^2)m^2s^2/\Lambda^6$.

We make a number of observations, arising from the above discussion, from the point of view of their future relevance:

- We note the existence of $m = 0$ limit in the expression (17).
- We note that the expression (17) can be expanded in powers of s at $m = 0$.
- We can suspect the exponential-like growth of (17) by an inspection of the integrand in (17).

3.2 Causality Violation at the One Loop level for the 2 Particle $\rightarrow$ 4 Particle process

To obtain the appropriate quantification for CV for this process in the one-loop approximation, we simplify the causality condition (6) using the unitarity condition (8). It reads,

$$\begin{aligned} H_2(x, y, z) = & iS_3(x, y, z) - iS_1(x)S_2(y, z) - iS_2(x, y)S_1(z) - iS_2(x, z)S_1(y) \\ & - iS_1(x)S_1(y)S_1(z) - iS_1(x)S_1(z)S_1(y) = 0 \end{aligned} \quad (22)$$

We find that for 2 particle $\rightarrow$ 4 particle process, the causality condition $H_2 = 0$ is violated by all the diagrams with *two* shadow propagators (see for example fig. 1(b)). To facilitate the calculation of CV in terms of the S-matrix elements, we use H_2 of (11). We find that H_2 consists of all diagrams for the 1-loop S-matrix with two barred lines. For concreteness, we shall first focus on the diagram of fig. 1(b). For this diagram the limit $m \rightarrow 0$ exists and in this limit the amplitude is given by⁴

$$\frac{27\lambda^3}{2\pi^2\Lambda^2} \int_0^1 d\tau_1 d\tau_2 \int_1^\infty \frac{d\tau_3}{(\tau_1 + \tau_2 + \tau_3)^2} \exp \left\{ \frac{k^2\tau_2}{\Lambda^2} + \frac{p^2\tau_3}{\Lambda^2} - \frac{(\tau_2 k + \tau_3 p)^2}{(\tau_1 + \tau_2 + \tau_3)\Lambda^2} \right\} \quad (23)$$

where $k = k_1 + k_2$ and $p = p_1 + p_2$. Other diagrams of this type can be now evaluated using the crossing symmetry. In this 6-point function no renormalization is required to be carried out. Therefore the leading order term, though a constant, will not be removed by renormalization. Leading order term (for $\frac{p^2}{\Lambda^2}, \frac{p \cdot k}{\Lambda^2}, \frac{k^2}{\Lambda^2} \ll 1$) is

$$\frac{27\lambda^3}{2\pi^2\Lambda^2} \int_0^1 d\tau_1 d\tau_2 \int_1^\infty \frac{d\tau_3}{(\tau_1 + \tau_2 + \tau_3)^2} = \frac{27\lambda^3}{2\pi^2\Lambda^2} (-4\ln 2 + 3\ln 3), \quad (24)$$

⁴After doing Schwinger parameterization and integrating over loop momentum

and since it is independent of any momenta this term remains the same for all other diagrams (total 15 in number). The *total* amplitude for the process is finite as $\Lambda \rightarrow \infty$ and behaves as (a quadratic function of momenta) $^{-1}$. Thus, the *relative* CV behaves as,

$$\frac{\text{a quadratic function of momenta}}{\Lambda^2}$$

and thus is expected to grow with energy.

On the other hand, for some of the various Lorentz invariants $\sim \Lambda^2$ (and having an appropriate sign), the truncation of the exponent as in (23) is not a good approximation. For example, for $k^2 = s \sim \Lambda^2$, $p^2 \sim \Lambda^2$ and $(p_3 + p_4)^2 \ll \Lambda^2$ and fixed, the expression (23) shows an exponential-like growth with s .

4 A general argument for $\phi\phi \rightarrow \phi\phi$

In this section, we shall attempt a simple way to generalize to all orders the result worked out for the $\phi\phi \rightarrow \phi\phi$ process in the previous section. For this purpose, we shall employ the results stated in the Appendix A regarding the infrared properties and analytic properties of the relevant amplitudes. The amplitude for the process and the CV in it is a Lorentz-invariant, dimensionless function only of the two independent Lorentz-invariants (say s and t) and of the parameters m and Λ . Let us parametrize the causality violating combination of S-matrix elements by $f(s, t, m^2, \Lambda^2)$. We shall first enumerate the properties expected of f :

- $f(s, t, m^2, \Lambda^2)$ vanishes as $\Lambda \rightarrow \infty$, i.e. in the local limit;
- $f(s, t, m^2, \Lambda^2)$ is dimensionless;
- $f(s, t, m^2, \Lambda^2)$ will have the crossing symmetry.

- $f(s, t, m^2, \Lambda^2)$ is finite at $m = 0$. We have given an argument regarding this in the Appendix A. To put the conclusions in it qualitatively, there is a cancellation of diagrams in the net CV amplitude; so that in the surviving diagrams, the infrared behavior is softer by the presence of (sufficiently many) barred propagators which have a better infrared behaviour. (Please note the barred propagator in section 2.1 for $m = 0$)⁵.
- $f(s, t, m^2, \Lambda^2)$ can be expanded in a series in s, t through $O(s^2)$ at $m = 0$. See Appendix A for the reasons.
- $f(s, t, m^2, \Lambda^2)$ is ambiguous upto momentum independent subtractions that vanish as $\Lambda \rightarrow \infty$.

In this case, the leading dependence will be a *linear* function of $\frac{s}{\Lambda^2}$ and $\frac{t}{\Lambda^2}$. The only function that also has the crossing symmetry is

$$f(s, t, m^2, \Lambda^2) = a \left\{ \frac{s}{\Lambda^2} + \frac{t}{\Lambda^2} + \frac{u}{\Lambda^2} \right\} + b \frac{m^2}{\Lambda^2} + O\left(\frac{1}{\Lambda^4}\right)$$

In view of $s + t + u = 4m^2$, we find that

$$f(s, t, m^2, \Lambda^2) = (b + 4a) \frac{m^2}{\Lambda^2} + O\left(\frac{1}{\Lambda^4}\right)$$

In view of the last observation regarding an ambiguity in the renormalization counterterms, we have,

$$f(s, t, m^2, \Lambda^2) = O\left(\frac{1}{\Lambda^4}\right)$$

⁵This and the following property can be addressed to *diagrammatically* by evaluating an arbitrary 1PI diagram and reducing it to the form of a Schwinger parametric integral such as of the form (23). The presence of a shadow propagator restricts the corresponding Schwinger parameter τ_i to $[0, 1]$ whereas the infrared divergence comes from $\tau_i \rightarrow \infty$. The property below about the expandability in s, t requires the study of the coefficients of s and t in the exponent in the integral. Also note the observations made at the end of the section 3.1.

modulo momentum-independent terms $O\left(\frac{1}{\Lambda^2}\right)$. Thus, the leading momentum-dependent CV in this process is of $O[\Lambda^{-4}]$ i.e. two powers of Λ suppressed as compared to what one would expect.

5 A possible scenario for escape from large scale causality violation at high energy scales

The scale Λ of the non-local standard model is, in principle, available phenomenologically from various precision experiments such as $(g-2)$ of the muon [12], the precision tests of standard model [20] etc, i.e. from experiments conducted at energies $\ll \Lambda$. As we had seen in sections 3 and 4, the causality violation can begin to grow very rapidly with the energy scale as the latter becomes a substantial fraction of the mass-scale Λ of the theory, so determined, and grows exponentially beyond it. It is possible that such a causality violation is actually present at high enough energies; in which case this will vindicate the use of non-local theory as the correct formulation of the physical theory, since such a causality violation cannot be explained by a LQFT. On the other hand, however, it is also possible that experiments detect no significant violation of causality even at such energies. One need not, however, necessarily conclude from it that the use of the non-local field theory to represent fundamental physics need be abandoned. In this section, we shall present a possible scenario that can provide an escape from such a predicament. In fact, we shall argue that such a predicament is indicative of a presence of a finer structure underlying the present theory.

We shall find it useful to employ the scenario presented in [15] (which we have briefly reviewed in section 2.4) for the present purpose also. In that work, we had argued that the condition $0 < Z < 1$ that is supposed to be satisfied by the

wave-function renormalization should in fact be taken literally in perturbation theory and can be implemented within the context of the non-local field theory. It can be used to predict the scale Λ that is present in the non-local formulation by a condition of the type

$$\frac{g^2}{16\pi^2} \ln \frac{\Lambda^2}{m^2} \lesssim 1.$$

[A similar conclusion was drawn in [14] from a discussion regarding renormalization program in a renormalizable field theory (including the standard model) where a condition $0 < Z < 1$ may not be derivable]. It was further proposed that the non-local field theory is an effective field theory and the effective field theory should be abandoned beyond energy scale $\gtrsim \Lambda$ where it must be replaced by a more fundamental theory with another coupling g' and mass scale Λ' . Such a scenario can also protect a non-local theory from a "causality-catastrophe".

We propose that should a large causality violation be unobserved at energies $\sim \Lambda$ (obtained from the above condition), it can be understood by invoking the above picture. In this picture, the calculations done in the non-local field theory at energy scales $\sim \Lambda$ are no longer very accurate, as the substructure of the theory becomes important. At these energy scales, the non-local field theory, which is being looked upon as an *effective* field theory, should be replaced by another non-local theory of its fundamental constituents having a new scale $\Lambda' \gg \Lambda$ and a new coupling constant. It is viable that in this new setting the causality violation may turn out to be in fact small as the energies are $\ll \Lambda'$.

In other words, in this view, the *calculated* large causality violation in NLQFT at large energy scales is argued as an artifact of the approximation that replaces the underlying theory of fundamental constituents by an effective non-local field theory.

The experimental observations regarding the CV at high enough energies may, in fact, enable one to distinguish between these two views regarding a

non-local theory: (i) A *fundamental* theory with a mass parameter Λ , (ii) an effective field theory valid upto a scale Λ .

6 Appendix A

In this appendix, we shall briefly deal with the infrared properties of the causality violating amplitudes constructed out of $H_n(y, x_1, \dots, x_n)$ that we have discussed in section 2.2 (see Eq.(4)). We have employed these properties in the section 4. On physical grounds, we can expect that quantities that characterize causality violation such as these should not have a sensitive dependence on m as $m \rightarrow 0$, that is, an absence of a mass singularity. We have already seen examples of lack of mass singularities in the calculations we have done. For example, in section 3.1, we have calculated the one loop amplitude arising from H_1 and found it to have

- Contribution from diagrams necessarily having one barred line (which have softer infrared properties);
- No singularity at $m = 0$ (see section 3.1.1);
- The amplitude is *analytic* (in s, t, u) at $m = 0$.
- There is no imaginary part, as there are no physical (i.e. having at least two smeared lines) intermediate states.

Further in section 3.2, we have seen the evaluation of the 1-loop amplitude based on H_2 . We found there that

- the contribution came from graphs with *at least two* barred lines;
- there is no singularity at $m = 0$;

- The amplitude is *analytic* in the quadratic Lorentz invariants at $m = 0$.

There is no imaginary part, as there are no physical intermediate states.

We draw attention to the fact that the H_n 's contain *specific* combinations of several terms involving various S_m 's. In H_n 's, there is a cancellation of diagrams: In the 4-point function of H_1 , the diagrams with only smeared propagator have cancelled out, while in the 6-point function of H_2 , diagrams with only smeared lines and those with *one* barred line have cancelled out. We shall argue that while each term in H_n may have infrared divergences/mass singularities as $m \rightarrow 0$, the combinations above cannot.

To see the argument, we consider the following quantity,

$$S[g(x), m]S^\dagger[g(x), m] \equiv 1 \quad \text{for all } m \text{ including } m \rightarrow 0$$

We note that the left hand side has a smooth limit as $m \rightarrow 0$ and consequently has no mass singularities in its matrix elements. We now imagine a small *local* variation in $g(x): g(x) \rightarrow g(x) + \delta g(x)$ in the first factor. We argue that *local* variations such as these cannot affect the infrared properties of the theory, which necessarily arise from large distances. Thus, the following operator: $S[g(x) + \delta g(x), m]S^\dagger[g(x), m]$ or equivalently the operator

$$O(y) \equiv i \frac{\delta S[g, m]}{\delta g(y)} S^\dagger[g, m]$$

has a smooth limit as $m \rightarrow 0$, i.e. has no mass singularity in its matrix elements.

We note the $H_n(y, x_1, \dots, x_n)$ has been defined as,

$$H_n(y, x_1, \dots, x_n) \equiv \frac{\delta^n}{\delta g(x_1) \delta g(x_2) \dots \delta g(x_n)} O(y) \Big|_{g=0}$$

and the causality violating combinations have been constructed out of these.

Next, we shall briefly deal with the analytic nature of the diagrams. It turns out that we can get at some of the analytic properties *without* detailed analysis of diagrams. Consider

$$S[g, m]S^\dagger[g, m] = 1$$

We differentiate with respect to $g(y)$ to obtain,

$$\frac{\delta S[g, m]}{\delta g(y)}S^\dagger[g, m] + S[g, m]\frac{\delta S^\dagger[g, m]}{\delta g(y)} = 0$$

Thus, $O(y) \equiv i\frac{\delta S[g, m]}{\delta g(y)}S^\dagger[g, m]$ is a hermitian operator [18]. Now consider the diagonal matrix elements of $O(x)$ between two 2-scalar states $|\alpha\rangle \equiv |k_1, k_2\rangle$. Then,

$$\langle \alpha | O(y) | \alpha \rangle = \langle \alpha | i\frac{\delta S[g, m]}{\delta g(y)}S^\dagger[g, m] | \alpha \rangle = a \text{ real quantity}$$

Now, matrix elements of H_n (analogues of (9) and (10)) are constructed out of

$$\frac{\delta}{\delta g(x_1)} \frac{\delta}{\delta g(x_2)} \dots \frac{\delta}{\delta g(x_n)} \langle \alpha | i\frac{\delta S[g, m]}{\delta g(y)}S^\dagger[g, m] | \alpha \rangle = a \text{ real quantity}$$

by real operations (see e.g. Eq.(10)). Thus, the quantity $\langle \alpha | H_n | \alpha \rangle$ does not develop imaginary part and the associated singularities due to contributions from physical intermediate states and thus cannot have singularities. This implies that the amplitudes so calculated $f(s, t, m, \Lambda)$ are free of singularities at $t = 0$ for any $m \geq 0$. This, in particular, rules out the terms of the type $\frac{s \ln s}{\Lambda^2}$ at $m = 0$. Crossing symmetry rules out terms such as $\frac{t \ln t}{\Lambda^2}$.

References

- [1] C. Itzykson, and J. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1985).
- [2] See, e.g. K. Namsrai, *Nonlocal Quantum Field Theory and Stochastic Quantum Mechanics* (D. Reidel Publishing company).
- [3] D. A. Eliezer, and R. P. Woodard, Phys.Rev.D 40, 465 (1989).
- [4] D. A. Eliezer, and R. P. Woodard, Nucl.Phys.B 325, 389 (1989).
- [5] G. Kleppe and R. P. Woodard, Nucl. Phys. B388, 81 (1992)
- [6] J.Moffat Phys. Rev. D41,1177(1990)
- [7] E. D. Evens et al, Phys Rev D43, 499 (1991)
- [8] G. Kleppe and R. P. Woodard, Ann. Phys. (N.Y.) 221, 106 (1993)
- [9] See e.g. N. J. Cornish Mod.Phys.Letts.A 7,1895 (1992); M.A.Clayton et al Int. Jour. Mod. Phys.A 9, 4549 (1994) and references therein; J.W. Moffat,hep-th/0207198 and references therein; as well as references 10-13 below.
- [10] G. Kleppe, Phys. Lett. B256, 431-436 (1991)
- [11] J. Paris, and W. Troost, Nucl. Phys. B482, 373-402 (1996).
- [12] S. D. Joglekar and G. Saini, Z. Phys. C.76, 343-353 (1997)
- [13] A. Basu, and S. D. Joglekar, J. Math. Phys. 41, 7206-7219 (2000); A. Basu, and S. D. Joglekar, Euro.Phys. Journal-direct C4 3, 1-20 (2001).
- [14] S.D.Joglekar, J. Phys. A34, 2765-2776 (2001)
- [15] S.D.Joglekar,Int.J.Mod. Phys.A 16, (2001).

- [16] N. J. Cornish, Int.J.Mod. Phys.A 7, 6121-6157 (1992).
- [17] N.N.Bogolibov and D.V.Shirkov, 'Introduction to Theory of Quantized Fields' (3^{rd} ed.), John Wiley(1980).
- [18] See pg. 200-220 of [17] .
- [19] G. C. Hegerfeldt, Phys. Rev. Lett. 72, 596 (1995) and the references therein;
G. C. Hegerfeldt, Annalen Phys. 7, 716-725 (1998).
- [20] A.Ayyer and S.D.Joglekar [unpublished calculations]